

DOCKET NO.: 215918US0



IN THE UNITED STATES PATENT & TRADEMARK OFFICE

IN RE APPLICATION OF:

YUKI MIYAZAWA ET AL

SERIAL NO: 09/987,549

FILED: NOVEMBER 15, 2001

:

: EXAMINER: PRYOR, A. N.

:

GROUP ART UNIT: 1616

FOR: TILLERING PROMOTER FOR PLANT AND METHOD FOR TILLERING
PROMOTION OF PLANT

CERTIFICATE OF TRANSLATION

COMMISSIONER OF PATENTS
ALEXANDRIA, VIRGINIA 22313

Sir:

Masao SHIMOKOSHI, the undersigned, a translator, residing at 356-46,
Obayashi, Koshigaya-shi, Saitama-ken, 343-0021 Japan

hereby states:

- (1) that I am fluent in both the Japanese and English languages;
- (2) that I translated the certified copy of the priority document filed in the instant application on November 15, 2001, of Japanese Patent Application No. 358705/2000 filed in Japan on November 27, 2000, from Japanese to English;
- (3) that the attached English translation is a true and correct translation of Japanese Patent Application No. 358705/2000, to the best of my knowledge and belief; and
- (4) that all statements made of my own knowledge are true and that all statements made on information and belief are believed to be true and further, that these statements are made with the knowledge that willful false statements and the like are punishable by fine or imprisonment, or both, under 18 USC 1001, and that such false statements may jeopardize the validity of the application or any patent issuing thereon.

Date: June 25, 2004

By:

Masao Shimokoshi

Masao SHIMOKOSHI

JAPAN PATENT OFFICE

This is to certify that the annexed is a true copy of the following application as filed with this Office

Date of Application: November 27, 2000

Application Number: JP2000-358705

Applicant(s): AJINOMOTO CO., INC.

Commissioner, Patent Office

Certification Number

Application Number JP2000-358705

[Document Name] REQUEST FOR PATENT

[Reference No.] MA43862

[Date of Filing] November 27, 2000

[Addressee] Commissioner, Patent Office

[Inventor]

[Address] c/o Ajinomoto Co., Inc., AminoScience
Laboratories, 1-1, Suzuki-cho, Kawasaki-ku,
Kawasaki-shi, Kanagawa-ken

[Name] Yuki MIYAZAWA

[Inventor]

[Address] c/o Ajinomoto Co., Inc., AminoScience
Laboratories, 1-1, Suzuki-cho, Kawasaki-ku,
Kawasaki-shi, Kanagawa-ken

[Name] Masahiko KURAUCHI

[Inventor]

[Address] c/o Ajinomoto Co., Inc., AminoScience
Laboratories, 1-1, Suzuki-cho, Kawasaki-ku,
Kawasaki-shi, Kanagawa-ken

[Name] Hiroyuki SATO

[Applicant for Patent]

[Code] 000000066

[Name] AJINOMOTO CO., INC.

[Agent]

[Code] 100064687

[Patent Attorney]

[Name] SHIMOKOSHI, Masao

[Phone No.] 03-5205-2384

Certification Number

Application Number JP2000-358705

[Agent]

[Code] 100102668

[Patent Attorney]

[Name] SAEKI, Norio

[Phone No.] 03-5205-2521

[Presentation of Fees]

[Deposit Book No.] 049401

[Amount of Payment] ¥21,000

[List of Attachments]

[Attachment Name] Specification 1

[Attachment Name] Abstract 1

[General Authorization] 9607453

[Proof] Requested

Certification Number

Application Number JP2000-358705

[Document Name] SPECIFICATION

[Title of the Invention] Tillering Promoter for Plant and
Method for Tillering Promotion of Plant

[Scope of Claim for Patent]

1. A tillering promoter for a plant comprising at least one amino acid selected from the group consisting of arginine, glutamine and proline as an effective ingredient.

2. The tillering promoter for a plant according to claim 1, wherein said amino acid is proline.

3. The tillering promoter for a plant according to claim 1 or 2, which contains also inosine in addition to said amino acid(s).

4. A tillering promoter for a plant, which is in the form of a kit comprising at least one amino acid selected from the group consisting of arginine, glutamine and proline to be applied onto plant leaf surfaces, and inosine to be applied to the subterranean part(s) of the plant.

5. The tillering promoter for the plant according to any one of claims 1 to 4, wherein said the plant is lawn grass.

6. The tillering promoter for a plant according to any one of claims 1 to 5, wherein the concentration of said amino acid(s) is in a range of 0.2 to 2,000 ppm.

7. The tillering promoter for a plant according to claim 6, wherein the concentration of said amino acid(s) is in a range of 100 to 300 ppm.

8. A method for tillering promotion of lawn grass wherein at least one amino acid selected from the group consisting of

Certification Number

Application Number JP2000-358705

arginine, glutamine and proline is applied onto lawn grass leaf surfaces, and inosine is applied to the subterranean part(s) of the lawn grass.

[Detailed Explanation of the Invention]

(0001)

(Technical Field Pertinent to the Invention)

The present invention relates to a tillering promoter for a plant comprising, as the effective ingredient(s), an amino acid, especially at least one of arginine, glutamine and proline which is effective for increasing tillering (as well known, branching out from the joints of stems near to the roots of rice, barley and wheat, lawn grass or the like) and weight of living plant, or inosine in addition to the amino(s) acid, and to the use of such a tillering promoter for a plant, or it relates to a method for tillering promotion of a plant wherein the amino acid(s) and inosine are each applied to the same part(s) or different part(s) of the plant at the same time or at slightly different times (i.e., used in combination).

(0002)

(Related Art)

The present inventors have previously developed a withering preventing and quick-acting nutrition supplementing agent for a gramineous plant comprising proline, one kind of amino acid, or inosine in addition thereto, as the effective ingredient(s) (Japanese Patent Application No. 308,281/1999). As a result of the studies for finding out further novel effects, the present inventors have achieved the present invention.

Certification Number

Application Number JP2000-358705

(0003)

Hitherto, some examples have been known wherein an amino acid-related compound, e.g., proline is applied to a plant.

(0004)

For example, (a) Japanese Patent Publication No. 42,566/1971 discloses a promoter of flower bud formation comprising at least one of uracil and cytosine, and proline.

(0005)

However, since the increase of the number of tillering is not observed in this promoter of flower bud formation and the method of application is different from the case of the present invention, the promoter is obviously different from the present invention.

(0006)

Furthermore, for the promotion of the tillering of lawn grass, there are known some examples wherein substances having indole skeleton(s) or plant hormone(s).

(0007)

For example, (a) Japanese Patent Application Laid-open (Kokai) No. 267,803/1995 discloses a tillering promoter for a gramineous plant comprising an fluorine-containing indolebutyric acid derivative, e.g., an alkyl ester of 4,4,4-trifluoro-3-(indole-3-)butyric acid.

(0008)

However, this tillering promoter is a chemical fertilizer and thus, it is obviously different from the present invention wherein amino acids, i.e., natural products are utilized, in

Application Number JP2000-358705

view of the influence against the environment.

(0009)

Moreover, (b) Japanese Patent Application Laid-Open (Kokai) No. 82,113/1995 discloses a growth promoter of lawn grass comprising a gibberellin and a cytokine.

(0010)

However, this growth promoter for lawn grass comprises plant hormones of gibberellin and cytokine, and thus, there is a possibility of exerting undesirable influences on the natural environment, so that the promoter is obviously different from the present invention wherein an amino acid is used.

(0011)

Furthermore, (c) Japanese Patent Application Laid-Open (Kokai) No. 201,914/1991 discloses a method for the promotion of the rooting and tillering of rice seedlings for transplantation, which comprises applying an amino acid fermentation liquid (containing, for example, proline, alanine, valine, and glutamic acid all together) during raising of seedlings in wet-rice farming.

(0012)

However, in this method of tillering promotion, proline is not solely used, and the amount of the amino acid fermentation liquid to be used is defined as total concentration of the amino acids, so that these amino acids are placed on the same level, i.e., are treated as equivalents.

(0013)

Incidentally, for example, lawns are utilized in many

Certification Number

Application Number JP2000-358705

places such as parks, gardens and ball game fields, and are particularly indispensable for golf courses. However, a large quantity of fertilizers and pesticides has hitherto been used for maintenance of lawn grass and the use has become a big problem in view of the environmental aspect.

(0014)

In particular, the stress by high temperature causes cold-district type lawn grass indirect growth stop and direct withering, but there is only a measure of good air-ventilation against the problems.

(0015)

(Problems to be Solved by the Invention)

Thus, it is an object of the present invention to provide a tillering promoter for a plant which enables the reduction of the amount of fertilizer hitherto used in a large amount and the maintenance of a satisfactory green state by promoting the tillering of a plant, especially lawn grass, without requiring the application of pesticides and plant hormones, also exhibits growth promotion and coloring promotion of leaves, is not a chemical fertilizer, and does not adversely affect the environment and men and beasts. Another object is to provide a method for applying the same.

(0016)

(Means for Solving the Problems)

As a result of extensive studies for achieving the aforementioned objects, the present inventors have for the first time found that an amino acid, especially proline, exhibits a

Certification Number

Application Number JP2000-358705

remarkable effect on the promotion of the tillering of lawn grass and inosine has an action of enhancing the tillering effect of the amino acid by promoting the growth of a plant. Accordingly, they have accomplished the present invention based on the findings.

(0017)

Namely, the present invention relates to a tillering promoter for a plant comprising an amino acid, especially proline, or inosine in addition thereto, as effective ingredient(s), and a method for tillering promotion for a plant wherein the tillering promoter is especially applied onto the leaf surfaces.

(0018)

(Embodiment of the Invention)

The following will explain the present invention in detail.

(0019)

The target plants to which the tillering promoter of the present invention is to be applied include preferably lawn grass, but also rice, fruits vegetables, leafy vegetables, decorative plants, and the like.

(0020)

As the amino acids, there may be mentioned arginine, glutamine and proline, and especially proline is effective, but these amino acids are not necessarily to be a purified one. They may be in the form of a protein hydrolysate or a mixture of amino acids containing a large quantity of proline unless it exerts an adverse effect. Each amino acid is preferably a product having

Certification Number

Application Number JP2000-358705

a high purity of 90% or more (ratio of the amino acid per the total solute excluding the inosine in the tillering promoter for a plant of the present invention to be applied onto leaf surfaces).

(0021)

The tillering promoter for a plant of the present invention comprising an amino acid, especially proline, as the effective ingredient can be prepared into a form wherein the effective ingredient is dissolved optionally in an appropriate solvent such as water or the like. Moreover, the promoter can be formulated into a powder, granules, or tablets by using an optional filler or binder. The application method is preferably foliar application. In this case, an amino acid concentration of 0.2 ppm to 0.2% (2,000 ppm), preferably 10 to 300 ppm, further preferably 100 to 300 ppm is effective. This is because no effect is exhibited at a concentration lower than the range, and there is a possibility of withering by excessive fertilization at a concentration higher than the range. By the way, in the case of dissolving in a solvent, it is optional to formulate the promoter by incorporating a fungicide, a surfactant, or a preservative in view of the prevention of rot. Furthermore, in the case of foliar application, the combined use with a spreader is effective.

(0022)

As the way of application of the tillering promoter including application timing, there may be mentioned application as an additional fertilizer, application after mowing grass,

Certification Number

Application Number JP2000-358705

or the like. With regard to the way of fertilizing, it is particularly effective to apply amino acid(s), especially proline, to the above-ground part(s) such as foliar application or the like, and inosine to subterranean part(s) by spraying to soil, addition to hydroponic medium, or the like.

(0023)

The foliar application of proline is effective not only for tillering promotion of a plant but also as a means for promoting the growth such as prevention of withering, feeding of fast-acting nitrogen, and the like. Moreover, the promotion of coloring leaves is also observed.

(0024)

Application amounts of the tillering promoter for a plant of the present invention vary depending on the application timing, the kind of plants, cultivation density, growing stage, and so on. In short, the amounts may be ones in which the tillering of the plant cultivated by using the tillering promoter of the present invention are superior to the tillering of the plant cultivated under entirely the same conditions with the exception that the tillering promoter of the present invention is not applied. It is possible to determine the amounts by some preliminary comparative test which is easy to carry out for those who are skilled in the art. For example, in the case of foliar application in a liquid form (as an agent for foliar application), proline may be applied at a low concentration, e.g., as low as 0.2 ppm within the above concentration range. That is, the tillering promotion of lawn grass is effected at such a low

Certification Number

Application Number JP2000-358705

concentration. In addition, the application amounts of inosine may be in a range of 0.05 to 1 ppm to soil (5 to 100 g per 100 tons of soil), and in the case of hydroponic cultivation, inosine may be applied in an amount of 0.1 to 2 ppm to hydroponic medium.

(0025)

By the way, the tillering promoter for a plant of the present invention may be in a form containing also (mixing) inosine in addition to a predetermined amino acid as explained in the above. However, the amino acid and inosine may be, of course, applied each at the same time or at slightly different times. Such an application way is also an embodiment of the present invention. As a formulation suitable for such an application way, there may be mentioned a kit form wherein the amino acid and inosine are packaged separately and both the packages are made one set.

(0026)

(Examples)

In the following will be explained the present invention in further detail with reference to the Examples.

(0027)

Example 1: Effect of proline on hydroponic cultivation of lawn grass (1)

Seedlings of grass (a European grass; bent grass) were raised and divided into four groups of A to D, and hydroponic cultivation was carried out (Table 1 shown below). Concerning Groups B and D, inosine was added to each hydroponic medium in such amount that the concentration became 2 ppm. Concerning Groups C and D, proline was applied onto leaf surfaces once a

Application Number JP2000-358705

week in an amount of 20 ppm. Group A was the control. Upon confirmation on the 40th day, the tillering promotion was clearly observed in the two groups of C and D wherein proline had been applied onto leaf surfaces, but no tillering promotion was observed in Groups A and B which had not been applied.

Furthermore, five average stocks which had not been withered were sampled from each group and examined. As shown in the following Table 1, with regard to all of the root length, leaf length and number (above-ground parts), total weight of the living plant, and tillering number, good growth was found in the proline-treated plots. Especially, it was confirmed that the effect became more remarkable when inosine was used in combination.

(0028)

Table 1: Comparison of the lawn grasses
between the groups (5 roots each)

	Average Root length (cm)	Average leaf length (cm)	Number of leaves per one root	Total weight of living plants per five roots (g)	Number of tillering per one root
A (Control)	1	18	5	0.28	1
B (Inosine)	2	20	7	0.40	2
C (Proline)	1	22	16	0.53	5
D (Inosine + Proline)	3	22	23	1.00	6

(0029)

Example 2: Effects of proline, glutamine, arginine and urea
on hydroponic cultivation of grass

Seedlings of grass (a European grass: bent grass) were raised and divided into five groups of A to E, and hydroponic cultivation was carried out (Table 2 shown below). Group A was

Application Number JP2000-358705

the control. Concerning Group B, proline was applied onto leaf surfaces once a week in an amount of 20 ppm. And, concerning the three groups of C to E, aqueous solutions of arginine, glutamine, and urea were applied onto leaf surfaces, respectively, once a week in such a amount that the nitrogen amount was the same as that of the proline in Group B. Upon confirmation on the 40th day, the tillering promotion was clearly observed in the groups of B to D, but no tillering promotion was observed in the groups of A and E.

(0030)

Furthermore, five average stocks which had not been withered were sampled from each group and examined. As shown in the following Table 2, with regard to the number of leaves and total weight of the living plant, good results were observed in the plots treated with proline, glutamine, and arginine, and the effects were more remarkable than that of urea which is a generally-used nitrogen source for foliar application.

(0031)

Table 2: Comparison of the lawn grasses
between the groups (5 roots each)

	Average root length (cm)	Average leaf length (cm)	Number of leaves per one root	Total weight of living plants per five roots (g)	Number of tillering per one root
A (Control)	2	14	8	0.33	2
B (Proline)	2	14	17	0.60	5
C (Arginine)	2	18	9	0.51	3
D (Glutamine)	2	18	11	0.66	3
E (Urea)	2	15	6	0.34	1

(0032)

Example 3: Effect of proline on hydroponic cultivation of grass

Application Number JP2000-358705

(2)

Seedlings of grass (a European grass: bent grass) were raised and divided into six groups of A to F, and hydroponic cultivation was carried out (Table 3 shown below). At that time, inosine was added to the hydroponic media for all the groups in such amount that the concentration became 2 ppm in each hydroponic medium. Group A was the control. In the groups of B to F, proline was applied onto leaf surfaces once a week with the concentration being changed stepwise. Upon confirmation on the 40th day, the tillering promotion was clearly observed in the groups of B to F, but no tillering promotion was observed in Group A. However, inhibition of the growth or withering was observed in Group F.

(0033)

Furthermore, five average stocks which had not been withered were sampled from each group and examined. As shown in the following Table 3, with regard to the number of leaves and total weight of the living plant, good results were observed in the plots treated with proline at a concentration of 2 ppm to 0.2%.

(0034)

Application Number JP2000-358705

Table 3: Comparison of the lawn grasses
between the groups (5 roots each)

	Average root length (cm)	Average leaf length (cm)	Number of leaves per one root	Total weight of living plants per five roots (g)	Number of tillering per one root
A (Control)	3	22	14	0.59	2
B (2 ppm Inosine)	4	24	14	1.04	3
C (20 ppm Proline)	4	24	15	0.95	4
D (200 ppm Proline)	5	24	24	1.01	4
E (0.2% Proline)	5	23	14	0.90	3
F (2.0% Proline)	3	19	11	0.40	3

(0035)

(Effects of the Invention)

According to the present invention, the application of at least one of proline, arginine and glutamine, or the application of inosine in addition to the amino acid(s) promotes the tillering of a plant, especially lawn grass, and also affords a nutrition effect easily.

Certification Number

Application Number JP2000-358705

[Document Name] ABSTRACT

[Abstract]

(Problems to be Solved by the Invention) To provide a tillering promoter for a plant which enables the reduction of the amount of fertilizer hitherto used in a large amount and the maintenance of a satisfactory green state by promoting the tillering of a plant, especially lawn grass, without requiring the application of pesticides and plant hormones, also exhibits growth promotion and coloring promotion of leaves, is not a chemical fertilizer, and does not adversely affect the environment and men and beasts.

(Means for Solving the Problems) A tillering promoter for a plant comprising an amino acid, especially proline, or inosine in addition thereto, as effective ingredient(s), and a method for tillering promotion for a plant wherein the tillering promoter is especially applied onto the leaf surfaces.

Certification Number